

Accessing COMETRIQ® (cabozantinib) Capsules

Specialty Pharmacies and Distributors

Contact one of these authorized in-network specialty pharmacies or distributors for access to COMETRIQ.

Specialty Pharmacies (US)	Phone	Fax	Website
Accredo Specialty Pharmacy	1-877-732-3431	1-866-352-3973	accredo.com
CVS Caremark Specialty Pharmacy	1-877-233-9920	1-855-296-0210	cvsspecialty.com
Optum Specialty Pharmacy	1-855-427-4682	1-877-342-4596	specialty.optumrx.com

340B, VA and NCI/NCCN Cancer Centers	Phone	Fax/Email	Website
Cardinal Health Specialty Distribution*	1-855-855-0708	GMB-SPD- CSORDERENTRY@cardinalhealth.com	specialtyonline.cardinalhealth.com orderexpress.cardinalhealth.com
McKesson Plasma and Biologics	1-877-625-2566	1-888-752-7626	connect.mckesson.com

*NCCN Comprehensive Cancer Center members only.



Exelixis Access Services® (EASE) is your resource for questions and needs related to financial assistance for COMETRIQ.



CALL: 1-844-900-EASE
(1-844-900-3273)



Monday to Friday
8:00 AM to 8:00 PM (ET)

For more information about accessing COMETRIQ, visit www.cometriq.com/hcp/access/.

Please see Important Safety Information on the next page and click [here](#) for the full Prescribing Information for COMETRIQ.



INDICATION

COMETRIQ® (cabozantinib) is indicated for the treatment of patients with progressive, metastatic medullary thyroid cancer (MTC).

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

The Full Prescribing Information for COMETRIQ includes Warnings and Precautions for:

- Perforations and Fistulas
- Hemorrhage
- Thrombotic Events
- Impaired Wound Healing
- Hypertension and Hypertensive Crisis
- Osteonecrosis of the Jaw
- Diarrhea
- Palmar-Plantar Erythrodysesthesia
- Proteinuria
- Reversible Posterior Leukoencephalopathy Syndrome
- Embryo-Fetal Toxicity
- Hypocalcemia

ADVERSE REACTIONS

The most common adverse reactions ($\geq 25\%$) are diarrhea, stomatitis, palmar-plantar erythrodysesthesia (PPE), decreased weight, decreased appetite, nausea, fatigue, oral pain, hair color changes, dysgeusia, hypertension, abdominal pain, and constipation.

The most common laboratory abnormalities ($\geq 25\%$) are increased AST, increased ALT, lymphopenia, increased ALP, hypocalcemia, hypoalbuminemia, neutropenia, thrombocytopenia, hypophosphatemia, and hyperbilirubinemia.

Please see full [Prescribing Information](#).

You are encouraged to report negative side effects of prescription drugs to the FDA.
Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.